

## LA-UR-14-27226

Approved for public release; distribution is unlimited.

Title: Investigation of light-matter interactions: Photoluminescence properties of individual quantum emitters

Author(s): Ma, Xuedan

Intended for: Progress meeting for a possible collaboration

Issued: 2014-10-22 (Rev.1) (Draft)

---

**Disclaimer:**

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by the Los Alamos National Security, LLC for the National Nuclear Security Administration of the U.S. Department of Energy under contract DE-AC52-06NA25396. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.




# Investigation of Light-Matter Interactions: Photoluminescence Properties of Individual Quantum Emitters

**Xuedan Ma**

**Center for Integrated Nanotechnologies, Materials Physics &  
Applications Division  
Los Alamos National Laboratory**

# Research Techniques

- PL/Raman/Rayleigh scattering measurements
- Laser: CW/pulsed, ultrafast
- cryogenic measurements
- E-beam/photo-lithography




Slide 2




# Mainly Studied Topics


II-VI semiconductor  
quantum dots (QDs)




Semiconductor  
nanotubes/nanowires





Single-walled carbon  
nanotubes (SWCNTs)



# QDs Coupled to Surface Plasmons




- enhanced PL intensity
- suppressed blinking
- gray states




- Lifetime decreases with  $d$ .
- Lifetime is dependent on dipole orientation.

# Semiconductor Nanotubes/Nanowires






ZnO NT

ZnO NT + Pt (inner)



Courtesy of Liang





# PL of Single-Walled Carbon Nanotubes

---

- Photon Statistics of Individual Carbon Nanotubes at Room Temperature
- Oxygen Doping Modifies Excitonic Fine Structures of Carbon Nanotubes
- Influence of Exciton Dimensionality on Spectral Diffusion of Single-Walled Carbon Nanotubes




# Photon Statistics of Individual Carbon Nanotubes at Room Temperature



# Why Photon Statistics?


photovoltaics and solar cells



*ACS Nano, 3, 3638–3648 (2009), Aydil et al.*

- multiexciton generation rate

quantum information





*A. Neuzner, MPQ  
Nature, 484, 195 (2012), Ritter et al.*


- single photon source

# Second-order Correlation Function $g^{(2)}$ :

## Hanbury-Brown-Twiss Setup



CW laser




Pulsed laser

$$(R = I_{center}/I_{side})$$


- $g^{(2)}$ : a measure of photon statistics.

# Photon Statistics of CNTs: State of the Art




4.2 K

*PRL 100, 217401 (2008), Hoegele et al.*



9 K

*Nano Lett. 12, 1934 (2012), Stefan et al.*



$$R < 0.5$$




photon antibunching

- Photon antibunching: photons are separated from each other.



# Photon Statistics of CNTs: Mechanism


Low T



localized exciton:

- exciton – exciton annihilation (EEA)

Room T



diffusing exciton:


- exciton – exciton annihilation (EEA)
- exciton diffusion



*How exciton diffusion and EEA affect photon statistics in 1D system?*

# Individual SWNT Imaging

Wide field images



Stable PL




PL blinking/fluctuation


- Homogeneous illumination of the whole SWNTs with expanded laser beam
- Beam size:  $\sim 60 \times 60 \text{ }\mu\text{m}$
- SWNT length:  $< 15 \text{ }\mu\text{m}$

# Individual SWNT Imaging

Wide field image




Perpendicular to the tube axis



width (@ $1/e^2$ )  
= 635 nm



- Fitted with a Gaussian function
- Diffraction limited

Along the tube axis





$L \sim 4.2 \mu\text{m}$

# Gaussian Laser Spot




- Fitted with a Gaussian function
- Diffraction limited



- Beam size independent of power.



# Excitation Power Dependent PL images




- 985.51 nm, 17.7 nm
- Fitted with a Lorentz function

- elongated PL image

# PL Intensity Profile



- width > point spread function
- due to exciton diffusion





- Profile width changes with excitation power.

# 1D Diffusion Equation


$$\frac{\partial n(x, t')}{\partial t'} = -n(x, t') + \underbrace{L_D^2}_{\text{diffusion length}} \frac{\partial^2 n(x, t')}{\partial x^2} - \underbrace{C}_{\text{exciton-exciton annihilation coefficient}} n(x, t')^2$$

diffusion length

exciton-exciton  
annihilation coefficient




Intensity profile




decay curve

# Fitting Intensity Profiles




- diffusion length  $L_D$
- EEA coefficient  $C$






# Exciton Diffusion $\leftrightarrow$ Exciton-Exciton Annihilation



- Larger diffusion length/rate  
→ less efficient exciton-exciton annihilation  
→ smaller  $C$

# Excitation Power Dependent $g^{(2)}$ Measurements




Area ratio:  

$$R = I_{center}/I_{side}$$

minimum antibunching degree  
 $R_0$

- $R$  increases with increasing excitation power.

## $R_0 \Leftrightarrow$ diffusion length $L_D$




# Minimum Antibunching Degree $R_0$


$$R_0 = Q_{2X} / (mQ_{1X}) + (m-1)/m$$

$Q_{2X}$ : emission efficiency of two excitons  
 $Q_{1X}$ : emission efficiency of one exciton  
 $m$ :  $d/L_D$

$$L_D > d \quad R_0 = Q_{2X} / (mQ_{1X})$$




$$L_D < d \quad R_0 = (m-1)/m$$




- Section of SWCNT behaviors like a QD.
- Excitons behave like independent emitters.

# $R_0 \Leftrightarrow$ Diffusion Length $L_D$




# Summary




$$\frac{\partial n(x, t')}{\partial t'} = -n(x, t') + L_D^2 \frac{\partial^2 n(x, t')}{\partial x^2} - C \cdot n(x, t')^2$$

Diffusion length  $L_D$  and EEA coefficient  $C$




minimum antibunching degree  $R_0$



$$R_0 = Q_{2X}/(mQ_{1X})$$

$$R_0 = (m-1)/m$$




UNCLASSIFIED



# Oxygen Doping Modifies Excitonic Fine Structures of Carbon Nanotube




# Photoluminescence of SWCNTs: Application





field-effect transistor

*Cao et al. Nature Mater. 2005, 4*




light emitting diode

*Wang et al. Nano Lett. 2011, 11*



sensor

*Kruss et al. J. Am. Chem. Soc. 2014*




biological imaging

*Welsher et al. Nature Nanotech. 2009*




# Photoluminescence of SWCNTs: Limitation

## External factors




- single layer structure
- structural defects
- adsorbate molecules/atoms

## Intrinsic factors



*Spataru et al. Phys. Rev. Lett. 2005*




*Kiowski et al. Phys. Rev. Lett. 2007*

- dark states below the first bright state


# Controlled Doping of SWCNTs

## oxygen doping




Ghosh et al. Science 2010

- additional bright, red-shifted peak




Miyauchi et al. Nature Photon. 2013

## diazonium salt doping




Piao et al. Nature Chem. 2013

# Individual SWCNT Imaging at Low Temperature




Wide field image




- diffraction-limited localized emission

# Individual Pristine SWCNT




- individual symmetric (88%) peak at  $\sim 1.25$  eV  $\rightarrow E_{11}$
- linewidth: 4.4 meV

# Individual Oxygen-Doped SWCNT




- spectral splitting of  $E_{11}$  peak  $\rightarrow$  2 to 3 peaks (70%)
- additional red-shifted peak  $\rightarrow E_{11}^*$

# Time Evolution of $E_{11}$ peak




- stable peak position and intensity




- spectral splitting

# Line Shape of $E_{11}^*$ Peak




- asymmetric peak with long tail at the low energy side (83%)



# Line Shape of $E_{11}^*$ Peak



Krummheuer et al. *Phys. Rev. B* 2002



Galland et al. *Phys. Rev. Lett.* 2008





localized exciton

- due to coupling between localized exciton and 1D phonon
- $E_{11}^*$  exciton is highly localized.




# Additional Peaks




- Both  $A^+$  and  $A^-$  are asymmetric with long tail at the low energy side.
- $A^+$  and  $A^-$  excitons are also localized.

# Quantum Chemistry Computations





- Length of tube: 8 nm
- l: along tube axis; d: perpendicular to tube axis
- Geometry optimization: semi-empirical method (AM1 Hamiltonian, MOPAC program)
- Oscillator strengths and transition densities: Time-dependent Density Functional Theory (Gaussian 09, STO-3G basis set, B3LYP functional).

# Comparison between Experiment and Simulation





# Exciton Transition Density




$E_{11}$


- symmetric lineshape  
→ delocalized exciton



$A^-$





$E_{11}^*$




$A^+$

- asymmetric lineshape  
→ localized exciton

# Intensity Correlation between $E_{11}$ and $E_{11}^*$ peaks




- PL intensities of  $E_{11}$  and  $E_{11}^*$  peaks are negatively correlated.






# Excitonic Fine Structures




- Oxygen doping introduces potential wells localizing excitons.
- Potential fluctuation leads to negative intensity correlation between  $E_{11}$  and  $E_{11}^*$  peaks.


# Summary




- spectral splitting of  $E_{11}$  peak



- asymmetric  $E_{11}^*$ ,  $A^+$ , and  $A^-$  peaks



- negative intensity correlation between  $E_{11}$  and  $E_{11}^*$  peaks



- excitonic fine structures of oxygen-doped SWCNTs




# Influence of Exciton Dimensionality on Spectral Diffusion of Single-Walled Carbon Nanotubes




# Spectral Diffusion of SWCNTs

Ideal spectra




Real spectra





- Quantum information processing requires spectral-diffusion free emission → physical mechanism behind spectral diffusion

# PL spectra of Individual SWCNTs at 4K





# Quantum-confined Stark effect (QCSE)

CdSe QDs  
(Bawendi, Science 97)




CdSe/CdS nanorods  
(Weller, PRL 2004)




CdSe/ZnS nanorods  
(Banin, Nano Lett. 2005)

# SWCNTs at Cryogenic Temperature

4 K



localized emission





complete antibunching

*Hoegeler et al. Phys. Rev. Lett. 2008*

- Excitons in SWCNTs at 4K is localized to **QD-like** excitonic states.  
→ sublinear correlation between linewidth and peak shift.

# PL spectra of Individual SWCNTs at RT

Exciton in SWCNT is **diffusive** at RT.  
Can we still observe QCSE?






# Exciton Dimensionality $\leftrightarrow$ Correlation Coefficient

4K  
exciton 0D




sublinear  
correlation



RT  
exciton 1D




Superlinear  
correlation





# Single Point Charge Model



For a SWCNT at low electric field, an energy shift of the binding energy ( $\Delta E$ ) caused by QCSE can be described by:

$$\Delta E = k_b \frac{(edF)^2}{E}$$

$e$ : electron charge     $d$ : tube diameter  
 $E$ : tube binding energy at zero field  
 $F$ : electric field induced by surface charges  
 $k_b$ : constant

If we use a Gaussian stochastic model to describe spectral diffusion behavior:

$$\langle \delta(\Delta E)(t) \delta(\Delta E)(0) \rangle = \sigma^2 e^{-t/t_c} \quad \sigma : \text{spectral broadening in meV}$$

At time  $t=0$  we obtain

$$\langle \delta^2(\Delta E)(0) \rangle = \sigma^2$$

In a simple assumption, we assume that there is an external point charge located on the surface of a SWCNT that creates a local electric field:

$$F = \frac{q}{4\pi\epsilon r^2}$$

$\epsilon$ : permittivity of the surrounding environment  
 $r$ : exciton-charge distance

# Single Point Charge Model



At cryogenic temperature:

- Excitons are localized to quantum-dot-like states.
- Exciton-charge distance  $r$  and local environment permittivity  $\epsilon$ : constants.
- Exciton binding energy fluctuation is mainly induced by fluctuations in surface charges:

$$\delta F = \frac{1}{4\pi\epsilon r^2} \delta q$$



$$\sigma \propto \sqrt{\Delta E}$$





# Single Point Charge Model



At room temperature:

- Excitons are diffusive ( $\sim 100$  nm).
- Large fluctuation in exciton-charge distance  $r$ .
- Contributions of surface charge/dielectric environment fluctuation become negligible.



$$\delta F = -\frac{2q}{4\pi\epsilon r^3} \delta r$$



$$\sigma \propto \Delta E^{5/4}$$





# SWCNTs Coupled to Surface Plasmons




$$\sigma \propto \sqrt{\Delta E}$$




exciton “localization”  
at room temperature




# Numerical Simulations




- highly confined excitation regime ( $< 30\text{nm}$ )




- strongly enhanced decay rates in the gap  $\rightarrow$  chance of exciton diffusion greatly decreased.




# Summary




$$\sigma \propto \sqrt{\Delta E}$$






exciton: 0D




$$\sigma \propto \Delta E^{5/4}$$



exciton: 1D




$\sigma \propto \sqrt{\Delta E}$   
exciton: 1D



# Current Project

Realization of Telecom Range (1300 - 1550 nm) Single Photon Source from SWCNTs at Room Temperature

- Air stable oxygen-doped SWCNTs
- Photon statistics of 1100 nm and 1300 nm peaks



# Acknowledgements:

---



Han Htoon  
Stephen Doorn  
Sergei Tretiak  
Andrei Piryatinski  
Jinkyong Yoo  
Jared Crochet  
Benjamin Mangum  
Juan Duque  
Xiaoying Pang  
Feng Wang  
Oleksiy Roslyak  
Hisato Yamaguchi  
Lyudmyla Adamska



U.S. DEPARTMENT OF  
**ENERGY**

Office of  
Science



UNCLASSIFIED

Slide 56

Operated by Los Alamos National Security, LLC for the U.S. Department of Energy's NNSA



---

**Thanks for your attention!**